


Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: <http://www.pediatr-neonatal.com>

## LETTER TO THE EDITOR

# Role of Pregnancy-induced Cholestasis in the Relation Between Mothers' Lipid Profile and Neonates' Macrosomia



To the Editor,

We appreciate Sahoo et al.<sup>1</sup> for their attention to our paper<sup>2</sup> recently published in *Pediatrics and Neonatology*. Pregnancy-induced cholestasis (PIC) affects 0.05–4% of pregnant women.<sup>3</sup> Since our cases were healthy and asymptomatic for PIC, this percentage will decrease to a very small value. In addition, our study participants were not at risk for PIC because they had neither a history of liver disease nor a twin pregnancy.<sup>3,4</sup> Ruling out cases with preterm labor (a complication of PIC in babies),<sup>4</sup> diabetes mellitus or positive screening test for diabetes mellitus, or obese persons<sup>5</sup> will also decrease the chance of including pregnant women with PIC in our study. Therefore, in the 154 cases we studied, the prevalence of PIC is negligible and will not affect our linear regression model at all.

Moreover, confounder role of PIC and its association with triglyceride is under question or controversial.<sup>6,7</sup> By contrast, its association with pregnancy is weak when its prevalence is low in pregnant women.<sup>3</sup> Therefore, evaluating cases suspected to have PIC and excluding them from the study are not necessary.

## Conflicts of interest

The authors have no conflicts of interest relevant to this article.

## Acknowledgments

This project did not receive any financial support.

## References

1. Sahoo J, Mitra S, Nayak PK, Kamalanathan S. Comment on prediction of neonates' macrosomia with maternal lipid profile

of healthy mothers by Mossayebi et al. *Pediatr Neonatol* 2014; 55:497–8.

2. Mossayebi E, Arab Z, Rahmaniyan M, Almassinokiani F, Kabir A. Prediction of neonates' macrosomia with maternal lipid profile of healthy mothers. *Pediatr Neonatol* 2014;55:28–34.
3. Chappell LC, Williamson C. Liver: links between hepatobiliary diseases in and beyond pregnancy. *Nat Rev Gastroenterol Hepatol* 2013;10:387–8.
4. Šimják P, Pařízek A, Vitek L, Černý A, Adamcová K, Koucký M, et al. Fetal complications due to intrahepatic cholestasis of pregnancy. *J Perinat Med* 2015;43:133–9.
5. Martineau M, Raker C, Powrie R, Williamson C. Intrahepatic cholestasis of pregnancy is associated with an increased risk of gestational diabetes. *Eur J Obstet Gynecol Reprod Biol* 2014;176:80–5.
6. Dai H, Jin H, He J. The changes of lipid metabolism and hemorheology in intrahepatic cholestasis during pregnancy. *Zhonghua Fu Chan Ke Za Zhi* 1997;32:728–31 [Article in Chinese].
7. Zhang Y, Hu L, Cui Y, Qi Z, Huang X, Cai L, et al. Roles of PPAR $\gamma$ /NF- $\kappa$ B signaling pathway in the pathogenesis of intrahepatic cholestasis of pregnancy. *PLoS One* 2014;9:e87343.

Ali Kabir\*

Department of Epidemiology, Faculty of Public Health,  
Shahid Beheshti University of Medical Sciences,  
Tehran, Iran

Center for Educational Research in Medical Sciences,  
Tehran University of Medical Sciences, Tehran, Iran

Elaheh Mossayebi

Department of Obstetrics and Gynecology, Shahid  
Akbarabadi Hospital, Iran University of Medical Sciences,  
Tehran, Iran

\*Corresponding author. Nikan Health Researchers Institute,  
Unit 9, Number 1, 3rd floor, 3rd Bahar Alley, Ashrafi Isfahani  
Highway, Poonak Square, Tehran, Iran.  
E-mail address: [aikabir@yahoo.com](mailto:aikabir@yahoo.com) (A. Kabir)

Nov 10, 2014